

For the use of a Registered Medical Practitioner or Hospital or a Laboratory only

UNIDERM NM

1. Generic name

Clobetasol Propionate, Neomycin Sulphate and Miconazole Nitrate Cream

2. Qualitative and quantitative composition

Clobetasol Propionate I.P....0.05% w/w

Miconazole Nitrate I.P.....2% w/w

Neomycin Sulphate I.P.....0.5% w/w

Chlorocresol I.P.0.1% w/w

(As preservative)

In a cream base.....q.s.

The excipients used are Butyl Hydroxytoulene, Butyl Hydroxyanisole, Cetodet-500, Hard Paraffin Wax, Di-Sodium Edetate, Methyl Hydroxy Benzoate, Propy Hydroxy Benzoate, Propylene Glycol, Polysorbate-20, White Petrollium Jelly, Colour Quinilone Yellow, Bronopol, Colour Tartazine Supra, Sodium Hydroxide Pellets.

3. Dosage form and strength

Dosage form: Topical Cream

Strength: Clobetasol Propionate 0.5mg/g, Neomycin Sulphate I.P. 5mg/g, Miconazole Nitrate I.P. 20mg/g and Chlorocresol I.P. 1mg/g (as preservative).

4. Clinical particulars

4.1. Therapeutic indication

For use only in condition of severe Acute Uveitis in adults only.

Remark: FDC shall not be used continuously for more than one week without re-evaluation by the physician.

4.2. Posology and method of administration

Sufficient amount of cream should be applied to cover the affected areas twice daily (morning and evening). Rub the cream in gently and completely.

UNIDERM NM Cream shall not be used continuously for more than one week without re-evaluation by the physician. The total dosage should not exceed 50 grams per week because of the potential of the clobetasol propionate to suppress the hypothalamic-pituitary-adrenal (HPA) axis. Further, treatment beyond 2 consecutive weeks is not recommended. Repeated short courses may be used to control exacerbations.

Or, as prescribed by the physician.

Method of administration

For topical use in adults and children over 2 years of age.

4.3. Contraindications

UNIDERM NM Cream is contraindicated in following cases:

- Hypersensitivity to clobetasol or to neomycin or to miconazole or to any component of

the formulation.

- Rosacea.
- Acne vulgaris.
- Perioral dermatitis.
- Perianal and genital pruritus
- Primary cutaneous viral infections (e.g., herpes simplex, chickenpox).
- Otitis externa with a perforated eardrum (because of risk of ototoxicity).

4.4. Special warnings and precautions for use

For external use only.

UNIDERM NM Cream must not come in contact with the eyes. Also, it should not be used with occlusive dressings. Due to the known ototoxic and nephrotoxic potential of neomycin sulphate, the use of UNIDERM NM cream in large quantities, or on large areas for prolonged periods of time is not recommended in circumstances where significant systemic absorption may occur.

Clobetasol Propionate

Use of potent topical corticosteroids, including clobetasol propionate, should be avoided on areas with thin and sensitive skin, such as on the face, groin or in the skin folds. As with other highly active corticosteroids, therapy should be discontinued when control has been achieved. If no improvement is seen within 2 weeks, reassessment of diagnosis may be necessary. Long-term continuous therapy should be avoided. Because of a higher ratio of skin surface area to body mass, children are at a greater risk than adults for HPA axis suppression when they are treated with topical corticosteroids. They are therefore also at greater risk of adrenal insufficiency after withdrawal of treatment, and of Cushing's syndrome while on treatment. Adverse effects including striae have been reported with inappropriate use of topical corticosteroids in infants and children.

Neomycin Sulphate

Neomycin can induce permanent sensorineural hearing loss due to cochlear damage, mainly destruction of hair cells in the organ of Corti (spiral organ - for hearing). The risk of ototoxicity is greater with prolonged use. Neomycin sulphate may cause cutaneous sensitization. A precise incidence of hypersensitivity reactions (primarily skin rash) due to topical neomycin is not known. Discontinue promptly if sensitization or irritation occurs.

Serious adverse reactions including neurotoxicity, ototoxicity and nephrotoxicity have occurred in patients receiving systemic aminoglycoside therapy. Although these effects have not been reported following topical use of aminoglycosides, caution is advised when used concomitantly with systemic aminoglycosides.

Pseudomembranous Colitis: Pseudomembranous colitis has been reported with the use of antibiotics and may range in severity from mild to life-threatening. Therefore, it is important to consider its diagnosis in patients who develop diarrhoea during or after antibiotic use. Although this is less likely to occur with topically applied neomycin, if prolonged or significant diarrhoea occurs or the patient experiences abdominal cramps, treatment should be discontinued immediately, and the patient investigated further.

Miconazole Nitrate

Severe hypersensitivity reactions, including anaphylaxis and angioedema, have been reported during treatment with miconazole topical formulations. If a reaction suggesting hypersensitivity or irritation should occur, the treatment should be discontinued.

4.5. Drug interactions

Clobetasol Propionate

CYP3A4 inhibitor drugs such as ritonavir and itraconazole, when co-administered with corticosteroids have been shown to inhibit the metabolism of corticosteroids leading to increased systemic exposure. The extent to which this interaction is clinically relevant depends on the dose and route of administration of the corticosteroids and the potency of the cytochrome P450 enzyme (CYP) 3A4 inhibitor.

Neomycin Sulphate

Following significant systemic absorption, neomycin sulphate can intensify and prolong the respiratory depressant effects of neuromuscular blocking agents. However, if used in accordance with the recommendations, systemic exposure to neomycin is expected to be minimal and drug interactions are unlikely to be significant. Concurrent use with other potentially nephrotoxic or ototoxic drugs should be avoided unless considered essential by the physician.

Miconazole Nitrate

Miconazole administered systemically is known to inhibit CYP3A4/2C9. Due to the limited systemic availability after topical application, clinically relevant interactions are rare. However, in patients on oral anticoagulants, such as warfarin, caution should be exercised, and anticoagulant effect should be monitored.

4.6. Use in special populations (such as Fertility, pregnancy, and lactation)

Pregnancy

Safety for use of UNIDERM NM Cream in pregnancy has not been established. It should only be used during pregnancy when considered mandatory by the physician, after careful assessment of the potential risks to the fetus.

Lactation

It is not known whether the components of UNIDERM NM Cream excrete in the breast milk after topical use. Nevertheless, caution should be exercised when this medication is administered to lactating women.

Paediatric Patients

Safety and effectiveness of this formulation has not been established in children below 2 years of age. Thus, due to safety concerns, UNIDERM NM Cream is not recommended in paediatric patients aged less than 2 years.

Geriatric Patients

Generally, no adjustment of dosage is required in the geriatric population. However, greater sensitivity of some older individuals cannot be ruled out.

4.7. Effects on ability to drive and use machines.

There have been no studies to investigate the effect of UNIDERM NM Cream on driving performance or the ability to operate machinery. From the adverse reaction profile of this product, effect on ability to drive and use machines is not expected with topical use of UNIDERM NM Cream.

4.8. Undesirable effects

UNIDERM NM cream is generally well tolerated. Following side effects may occur

occasionally with the individual components of UNIDERM NM Cream.

Clobetasol Propionate

The following adverse reactions have been reported with use of clobetasol propionate. The frequency of these adverse events is unknown.

Immune System Disorders - Hypersensitivity:

Local hypersensitivity reactions such as erythema, rash, pruritus, urticaria and allergic contact dermatitis may occur at the site of application and may resemble symptoms of the condition under treatment. If signs of hypersensitivity appear, application should be stopped immediately.

Endocrine Disorders - Features of Cushing's Syndrome:

As with other topical corticosteroids, prolonged use especially of large amounts, or treatment of extensive areas can lead to sufficient systemic absorption to produce the features of Cushing's syndrome. This effect is more likely to occur in infants and children, and if occlusive dressings are used. In infants, the nappy may act as an occlusive dressing. Provided the weekly dosage is less than 50 gram in adults, any suppression of the HPA axis is likely to be transient with a rapid return to normal values once the short course of steroid therapy has ceased. The same applies to children given proportionate dosage.

Vascular Disorders - Dilatation of the Superficial Blood Vessels:

Prolonged and intensive treatment with potent corticosteroid preparations may cause dilatation of the superficial blood vessels, particularly when occlusive dressings are used, or when skin folds are involved.

Skin and Subcutaneous Tissue Disorders:

Local skin burning, local atrophy, striae, thinning, pigmentation changes, hypertrichosis, exacerbation of underlying symptoms, pustular psoriasis. Prolonged and intensive treatment with potent corticosteroid (clobetasol) preparations may cause local atrophic changes, such as thinning and striae.

Neomycin Sulphate

Neomycin occasionally causes skin sensitization. Ototoxicity and nephrotoxicity have also been reported with use of neomycin in large quantities and/or for prolonged periods.

Miconazole Nitrate

Local side effects such as itching, burning, rash, and contact dermatitis have been reported with topical miconazole therapy.

Reporting of suspected adverse reactions

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via any point of contact of Torrent Pharma available at:
https://www.torrentpharma.com/index.php/site/info/adverse_event_reporting

By reporting side effects, you can help provide more information on the safety of this medicine.

4.9. Overdose

Clobetasol Propionate

Acute overdose is very unlikely to occur, however, in the case of chronic overdose or misuse,

the features of hypercortisolism may appear and in this situation topical steroids should be reduced or discontinued gradually, under medical supervision.

Neomycin Sulphate

If systemic absorption of neomycin sulphate is suspected, use of the product should be stopped and the patient's general status, hearing acuity, renal and neuromuscular functions should be monitored. Blood levels of neomycin sulphate should also be determined. Haemodialysis may reduce the serum level of neomycin sulphate.

Miconazole Nitrate

Excessive cutaneous use can result in skin irritation, which usually disappears after discontinuation of therapy. If accidental ingestion of large quantities of the product occurs, an appropriate method of gastric emptying may be used if considered necessary.

5. Pharmacological properties

5.1. Mechanism of Action

Clobetasol Propionate

The mechanism of anti-inflammatory activity of the topical corticosteroids involves induction of phospholipase A2 inhibitory proteins, collectively called lipocortins. It is postulated that these proteins control the biosynthesis of potent mediators of inflammation such as prostaglandins and leukotrienes by inhibiting the release of their common precursor, arachidonic acid. Arachidonic acid is released from membrane phospholipids by phospholipase A2.

Clobetasol propionate also produces anti-inflammatory response, partially due to vasoconstriction and decrease in collagen synthesis.

Neomycin Sulphate

Neomycin exerts its bactericidal effect by interfering with the protein synthesis of susceptible organisms. Neomycin is rapidly acting bactericidal agent; after diffusing through the bacterial cell membrane, it binds polysomes to affect protein synthesis. Neomycin disrupts the normal cycle of ribosomal function by interfering, at least in part, with the first step of protein synthesis. Neomycin also causes a misreading of the genetic code of the mRNA template, and this causes incorrect amino acids to be incorporated into the growing polypeptide chain, producing nonsense proteins.

Miconazole Nitrate

Miconazole is an imidazole antifungal agent that acts by interfering with the permeability of the fungal cell membrane. Miconazole inhibits biosynthesis of ergosterol, damaging the fungal cell wall, which increases permeability causing leakage of nutrients.

5.2. Pharmacodynamic Properties

Clobetasol Propionate

Like other topical corticosteroids, clobetasol propionate has anti-inflammatory, antipruritic, and vasoconstrictive properties. The major effect of clobetasol propionate on skin is a non-specific anti-inflammatory response, partially due to vasoconstriction and decrease in collagen synthesis.

Neomycin Sulphate

Neomycin sulphate is bactericidal against a wide range of Gram-positive and Gram-negative

bacterial pathogens including *Staphylococci*, *Streptococci*, *Escherichia*, *Enterobacter*, *Klebsiella*, *Hemophilus*, *Proteus*, *Salmonella* and *Shigella* species. It is also active against some strains of *Pseudomonas aeruginosa*, and against *Mycobacterium tuberculosis* and *Neisseria gonorrhoea*.

Miconazole Nitrate

Miconazole possesses a wide antifungal spectrum and has some antibacterial activity. Miconazole inhibits the growth of the common dermatophytes such as *Trichophyton rubrum*, *Trichophyton mentagrophytes*, and *Epidermophyton floccosum*, and the yeast-like fungi such as *Candida albicans*. Also, many Gram-positive bacteria including most strains of *Streptococcus* and *Staphylococcus* are susceptible to miconazole.

5.3. Pharmacokinetic Properties

Clobetasol Propionate

Percutaneous penetration of clobetasol propionate varies among individuals and can be increased by the use of occlusive dressings, or when the skin is inflamed or diseased. Following percutaneous absorption of clobetasol propionate, the drug probably follows the metabolic pathway of systemically administered corticosteroids i.e., metabolized primarily by the liver and then excreted by the kidneys. However, systemic metabolism of clobetasol has never been fully characterised or quantified.

Neomycin Sulphate

Although not absorbed through intact skin, topical neomycin is readily absorbed from large denuded, burned, or granulating areas. Greater and more rapid absorption occurs with neomycin cream than with the ointment. Plasma concentrations following topical application to open wounds, burns, or granulating surfaces are comparable to, or higher than, those achieved following oral preparation. Once neomycin is absorbed, it is rapidly excreted by the kidneys in active form. It has been reported to have a half-life of 2 to 3 hours.

Miconazole Nitrate

There is little absorption through skin or mucous membranes when miconazole nitrate is applied topically. Absorbed miconazole is bound to plasma proteins (88.2%) and red blood cells (10.6%). The small amount of miconazole that is absorbed is eliminated predominantly in faeces as both unchanged drug and metabolites.

6. Nonclinical properties

6.1. Animal Toxicology or Pharmacology

In fertility studies, subcutaneous administration of clobetasol propionate to rats at doses of 6.25 to 50 micrograms/kg/day produced no effects on mating, and fertility was only decreased at 50 micrograms/kg/day.


Subcutaneous administration of clobetasol propionate to mice (≥ 100 micrograms/kg/day), rats (400 micrograms/kg/day) or rabbits (1 to 10 micrograms/kg/day) during pregnancy produced foetal abnormalities including cleft palate.

In the rat study, where some animals were allowed to litter, developmental delay was observed in the F1 generation at ≥ 100 micrograms/kg/day and survival was reduced at 400 micrograms/kg/day. No treatment-related effects were observed in F1 reproductive performance or in the F2 generation.

7. Description

Clobetasol Propionate

Clobetasol Propionate is [(8S,9R,10S,11S,13S,14S,16S,17R)-17-(2-chloroacetyl)-9-fluoro-11-hydroxy-10,13,16-trimethyl-3-oxo-6,7,8,11,12,14,15,16-octahydrocyclopenta[a]phenanthrene -17-yl] propanoate. The empiric formula of $C_{25}H_{32}ClFO_5$ and its molecular weight is 467.0 g/mol. Its structural formula is:




Neomycin Sulphate

Neomycin Sulphate is (2R,3S,4R,5R,6R)-5-amino-2-(aminomethyl)-6-[(1R,2R,3S,4R,6S)-4,6-diamino-2-[(2S,3R,4S,5R)-4-[(2R,3R,4R,5S,6S)-3-amino-6-(aminomethyl)-4,5 dihydroxyoxan -2-yl] oxy-3-hydroxy-5-(hydroxymethyl) oxolan-2-yl]oxy-3 hydroxy cyclohexyl]oxyoxane-3,4-diol;methane. The empiric formula of $C_{24}H_{50}N_6O_{13}$ and its molecular weight is 630.7 g/mol. Its structural formula is:



Miconazole Nitrate

Miconazole Nitrate is 1-[2-(2,4-dichlorophenyl)-2-[(2,4-dichlorophenyl) methoxy]ethyl]imidazole; nitric acid. The empiric formula of $C_{18}H_{15}Cl_4N_3O_4$ and its molecular weight is 479.1 g/mol. Its structural formula is:



UNIDERM NM is yellow coloured soft mass. The excipients used are Butyl Hydroxytoulene, Butyl Hydroxyanisole, Cetodet-500, Hard Paraffin Wax, Di-Sodium Edetate, Methyl Hydroxy Benzoate, Propy Hydroxy Benzoate, Propylene Glycol, Polysorbate-20, White Petrollium Jelly, Colour Quinilone Yellow, Bronopol, Colour Tartazine Supra, Sodium Hydroxide Pellets.

8. Pharmaceutical particulars:

8.1. Incompatibilities:

Not applicable.

8.2. Shelf-life:

Do not use later than date of expiry.

8.3. Packaging information:

UNIDERM NM are packed in 15 gm tube.

8.4. Storage and handing instructions:

Stores protect from light, at a temperature not exceeding 25°C. Do not Freeze.

Avoid direct contact with eyes.

Keep the medicine out of reach of children.

Keep the tube tightly closed after use.

9. Patient Counselling Information

Ask the patients to inform the treating physicians in case of any of the below:

- Have any allergies.
- Have kidney or liver problems.
- Are pregnant or plan to become pregnant.
- Are breastfeeding or plan to breastfeed.
- Have any serious illness.
- Are taking any medicines (prescription, over the counter, vitamins, or herbal products)

10. Details of manufacturer

Manufactured by:

East African (India) Overseas

Plot No. 8, Pharmacity, Selaqui,

Dehradun-248 011 (Uttarakhand).

11. Details of permission or license number with date

Torrent Pharmaceuticals Ltd.,

Mfg Lic No. 94/UA/SC/P-2009 issued on 20.03.2017.

12. Date of revision

NA

MARKETED BY



TORRENT PHARMACEUTICALS LTD.

Indrad-382 721, Dist. Mehsana, INDIA.

IN/UNIDERM NM/Mar 2025/01/PI