

FLAVOSPAS TAB

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory Only

Abbreviated Prescribing information for FLAVOSPAS TAB [Flavoxate Hydrochloride 200 mg]

[Please refer the complete prescribing information available at www.torrentpharma.com]

PHARMACOLOGICAL PROPERTIES:

MECHANISM OF ACTION: The mechanism of action involves intracellular cyclic AMP accumulation and calcium blocking activity. It inhibits bladder contractions induced by various agonists or by electrical stimulation and inhibits the frequency of bladder voiding contractions. It increases bladder volume capacity, reduces the threshold and micturition pressure.

INDICATIONS: Flavoxate is indicated for the symptomatic relief of dysuria, urgency, nocturia, vesical supra-pubic pain, frequency and incontinence as may occur in cystitis, prostatitis, urethritis, urethro-cystitis and urethrotrigonitis. In addition, the preparation is indicated for the relief of vesico-urethral spasms due to catheterisation, cystoscopy or indwelling catheters; prior to cystoscopy or catheterisation; sequelae of surgical intervention of the lower urinary tract.

DOSAGE AND ADMINISTRATION: The recommended adult dosage is one tablet three times a day for as long as required. Flavoxate tablets are not recommended for children under 12 years of age.

CONTRAINDICATION: The following obstructive conditions: pyloric or duodenal obstruction, obstructive intestinal lesions or ileus, achalasia, gastrointestinal haemorrhage and obstructive uropathies of the lower urinary tract. Hypersensitivity to flavoxate hydrochloride or to any of the excipients of Flavoxate 200.

WARNINGS & PRECAUTIONS: Flavoxate should be used with caution in patients with suspected glaucoma, especially closed angle cases and in patients with serious, uncontrolled, obstructive disorders of the lower urinary tract. Where evidence of urinary infection is present, appropriate anti-infective therapy should be instituted concomitantly.

DRUG INTERACTIONS: N/A

ADVERSE REACTIONS: Eosinophilia, leukopenia, Hypersensitivity reactions, Drowsiness, dizziness, headache, mental confusion (especially in the elderly), nervousness, vertigo, Blurred vision, disturbances in eye accommodation, increased ocular tension, Palpitations, tachycardia, Diarrhoea, dry mouth, dyspepsia, dysphagia, nausea, vomiting, Angioedema, urticaria, erythema, rash, pruritis, Dysuria, Fatigue, hyperpyrexia.

MARKETED BY:



Torrent Pharmaceuticals Limited.

(Additional information is available on request)