

## Fungicide & SPOO-KT

**For the use of a Registered Medical Practitioner or a Hospital or a Laboratory Only**

Abbreviated Prescribing information for Fungicide & SPOO-KT [Ketoconazole 2.00 % + Zinc Pyrithione 1.00 %]  
[Please refer the complete prescribing information available at [www.torrentpharma.com](http://www.torrentpharma.com)]

### PHARMACOLOGICAL PROPERTIES:

**MECHANISM OF ACTION:** Ketoconazole is a synthetic derivative of imidazole-dioxolane, providing fungicidal and fungistatic activity. Zinc pyrithione suspension has antimicrobial and antifungal activity.

**INDICATIONS:** For the treatment of pityriasis and dandruff of the scalp.

**DOSAGE AND ADMINISTRATION:** As directed by the physician.

**CONTRAINDICATION:** Hypersensitivity to Ketoconazole, Zinc Pyrithione and other excipients.

**WARNINGS & PRECAUTIONS:** In patients who have been on prolonged treatment with topical corticosteroids, it is recommended that the steroid therapy be gradually withdrawn over a period of 2 to 3 weeks, while using Ketoconazole with Zinc Pyrithione Lotion, to prevent any potential rebound effect.

Keep out of the reach of eyes. If the lotion gets into the eyes, immediately, the eyes should be rinsed off with water.

**DRUG INTERACTIONS:** When prescribed in patients that have been treated with topical glucocorticosteroids for a long time, treatment with the latter should be continued and they should be gradually withdrawn for 2-3 weeks (to avoid the withdrawal syndrome).

Taking into account lack of significant systemic absorption, interaction with other medicinal products is unlikely.

It is not recommended to use it concomitantly with dermatological preparations containing glucocorticosteroids.

**ADVERSE REACTIONS:** Hypersensitivity, Dysgeusia, Folliculitis, Eye irritation, Alopecia, dry skin, hair texture abnormal, rash, skin burning sensation, acne, dermatitis contact, skin disorder, skin exfoliation, angioedema, urticaria, hair colour changes, Application site erythema, application site pruritus, application site bleeding, application site discomfort, application site dryness, application site inflammation, application site irritation application site paraesthesia, and application site reaction.

**TORRENT**  
PHARMA

### MARKETED BY:

Torrent Pharmaceuticals Limited.

**INFUNGICIDE & SPOO-KT/JAN-2026/01/ABPI**

(Additional information is available on request)