

PRODUCT NAME :	Tolterodine Tartrate Extended-Release Capsules	COUNTRY : US	LOCATION : Indrad	Supersedes A/W No.:	8102981	V. No.: 01	
ITEM / PACK :	Outsert	NO. OF COLORS: 1	REMARK :				
DESIGN STYLE :	Front	PANTONE SHADE NOS.:	SUBSTRATE : 40 gm/2 Bible Paper				
CODE :	8106403	Black	Activities	Department	Name	Signature	Date
DIMENSIONS (MM) :	490 x 340		Prepared By	Pkg.Dev			
ART WORK SIZE :	S/S		Reviewed By	Pkg.Dev			
DATE :	03-01-2026		Reviewed By	Quality			
		Font Size 6.5 pt_Medi Font Size 10 pt	Approved By	Quality			

Note: Pharma code/ Bar code and adjacent text must be visible on folded leaflet.

These details can be moved by printed to arrange pharma code/ Bar code and adjacent text visible on folded leaflet.

HIGHLIGHTS OF PRESCRIBING INFORMATION
These highlights do not include all the information needed to use TOLTERODINE TARTRATE EXTENDED-RELEASE CAPSULES safely and effectively. See full prescribing information for TOLTERODINE TARTRATE EXTENDED-RELEASE CAPSULES.

TOLTERODINE TARTRATE extended-release capsules

For oral administration
Initial U.S. Approval: December 2000

-----INDICATIONS AND USAGE-----

Tolterodine tartrate extended-release capsules are an antimuscarinic indicated for the treatment of overactive bladder with symptoms of urge urinary incontinence, urgency, and frequency. (1)

-----DOSAGE AND ADMINISTRATION-----

- 4 mg capsules taken orally once daily with water and swallowed whole. (2, 1)
- 2 mg capsules taken orally once daily with water and swallowed whole in the presence of:
 - mild to moderate hepatic impairment (Child-Pugh class A or B) (2,2)
 - severe renal impairment [Creatinine Clearance (CrCl) 10 to 30 mL/min] (2,2)
 - drugs that are potent CYP3A4 inhibitors. (2,2)
- Tolterodine tartrate extended-release capsules are not recommended for use in patients with CrCl <10 mL/min. (2,2)
- Tolterodine tartrate extended-release capsules are not recommended for use in patients with severe hepatic impairment (Child-Pugh Class C). (2,2)

-----DOSAGE FORMS AND STRENGTHS-----

Capsules: 2 mg and 4 mg (3)

-----CONTRAINDICATIONS-----

- Tolterodine tartrate extended-release capsules are contraindicated in patients with urinary retention, gastric retention, or uncontrolled narrow-angle glaucoma. Tolterodine tartrate extended-release capsules are also contraindicated in patients with known hypersensitivity to the drug or its ingredients, or to fesoterodine fumarate extended-release tablets which, like tolterodine tartrate extended-release capsules, are metabolized to 5-hydroxymethyl tolterodine. (4)
- Warnings and Precautions**
 - Anaphylaxis and angioedema requiring hospitalization and emergency medical treatment have occurred with the first or subsequent doses of tolterodine tartrate extended-release capsules. (5,1)
 - Urinary Retention: use caution in patients with clinically significant bladder outflow obstruction because of the risk of urinary retention. (5,2)
 - Gastrointestinal Disorders: use caution in patients with gastrointestinal obstructive disorders or decreased gastrointestinal motility because of the risk of gastric retention. (5,3)
 - Controlled Narrow-Angle Glaucoma: use caution in patients being treated for narrow-angle glaucoma. (5,4)
 - Central Nervous System Effects: Somnolence has been reported with tolterodine tartrate extended-release capsules. Advise patients not to drive or operate heavy machinery until they know how tolterodine tartrate extended-release capsules affect them. (5,5)
 - Myasthenia Gravis: use caution in patients with myasthenia gravis. (5,8)
 - QT Prolongation: consider observations from the thorough QT study in clinical decisions to prescribe tolterodine tartrate extended-release capsules to patients with a known history of QT prolongation or to patients who are taking Class IA (e.g., quinidine, procainamide) or Class III (e.g., amiodarone, sotalol) antiarrhythmic medications. (5,9)

FULL PRESCRIBING INFORMATION:

CONTENTS*

1 INDICATIONS AND USAGE

- Dosing Information
- Dosage Adjustment in Specific Populations

2 DOSAGE AND ADMINISTRATION

- Dosing Information
- Dosage Adjustment in Specific Populations

3 NONCLINICAL TOXICOLOGY

- Carcinogenesis, Mutagenesis, Impairment of Fertility

4 CLINICAL STUDIES

5 MYASTHENIA GRAVIS

6 HOW SUPPLIED/STORAGE AND HANDLING

7 PATIENT COUNSELING INFORMATION

- Angioedema
- Urinary Retention
- Gastrointestinal Disorders
- Controlled Narrow-Angle Glaucoma
- Central Nervous System Effects
- Hepatic Impairment
- Renal Impairment
- Myasthenia Gravis
- Use in Patients with Congenital or Acquired QT Prolongation

8 ADVERSE REACTIONS

- Clinical Trials Experience
- Post-marketing Experience

9 DRUG INTERACTIONS

- Potent CYP2D6 Inhibitors
- Potent CYP3A4 Inhibitors
- Other Interactions
- Other Drugs Metabolized by Cytochrome P450 Isoenzymes
- Drug-Laboratory-Test Interactions
- Other Anticholinergics

10 USE IN SPECIFIC POPULATIONS

- Pregnancy
- Lactation
- Pediatric Use
- Geriatric Use
- Renal Impairment
- Hepatic Impairment

11 OVERDOSAGE

12 DESCRIPTION

capsules. Advise patients not to drive or operate heavy machinery until they know how tolterodine tartrate extended-release capsules affect them. (5,5)

TOLTERODINE TARTRATE extended-release capsules

For oral administration
Initial U.S. Approval: December 2000

-----INDICATIONS AND USAGE-----

Tolterodine tartrate extended-release capsules are an antimuscarinic indicated for the treatment of overactive bladder with symptoms of urge urinary incontinence, urgency, and frequency. (1)

-----DOSAGE AND ADMINISTRATION-----

- 4 mg capsules taken orally once daily with water and swallowed whole. (2, 1)
- 2 mg capsules taken orally once daily with water and swallowed whole in the presence of:
 - mild to moderate hepatic impairment (Child-Pugh class A or B) (2,2)
 - severe renal impairment [Creatinine Clearance (CrCl) 10 to 30 mL/min] (2,2)
 - drugs that are potent CYP3A4 inhibitors. (2,2)
- Tolterodine tartrate extended-release capsules are not recommended for use in patients with CrCl <10 mL/min. (2,2)
- Tolterodine tartrate extended-release capsules are not recommended for use in patients with severe hepatic impairment (Child-Pugh Class C). (2,2)

-----DOSAGE FORMS AND STRENGTHS-----

Capsules: 2 mg and 4 mg (3)

-----CONTRAINDICATIONS-----

- Tolterodine tartrate extended-release capsules are contraindicated in patients with urinary retention, gastric retention, or uncontrolled narrow-angle glaucoma. Tolterodine tartrate extended-release capsules are also contraindicated in patients with known hypersensitivity to the drug or its ingredients, or to fesoterodine fumarate extended-release tablets which, like tolterodine tartrate extended-release capsules, are metabolized to 5-hydroxymethyl tolterodine. (4)
- Warnings and Precautions**
 - Anaphylaxis and angioedema requiring hospitalization and emergency medical treatment have occurred with the first or subsequent doses of tolterodine tartrate extended-release capsules. (5,1)
 - Urinary Retention: use caution in patients with clinically significant bladder outflow obstruction because of the risk of urinary retention. (5,2)
 - Gastrointestinal Disorders: use caution in patients with gastrointestinal obstructive disorders or decreased gastrointestinal motility because of the risk of gastric retention. (5,3)
 - Controlled Narrow-Angle Glaucoma: use caution in patients being treated for narrow-angle glaucoma. (5,4)
 - Central Nervous System Effects: Somnolence has been reported with tolterodine tartrate extended-release capsules. Advise patients not to drive or operate heavy machinery until they know how tolterodine tartrate extended-release capsules affect them. (5,5)
 - Myasthenia Gravis: use caution in patients with myasthenia gravis. (5,8)
 - QT Prolongation: consider observations from the thorough QT study in clinical decisions to prescribe tolterodine tartrate extended-release capsules to patients with a known history of QT prolongation or to patients who are taking Class IA (e.g., quinidine, procainamide) or Class III (e.g., amiodarone, sotalol) antiarrhythmic medications. (5,9)

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Tolterodine tartrate extended-release capsules are indicated for the treatment of overactive bladder with symptoms of urge urinary incontinence, urgency, and frequency [see **CLINICAL STUDIES** (14)].

2 DOSAGE AND ADMINISTRATION

2.1 Dosing Information

The recommended dose of tolterodine tartrate extended-release capsules is 4 mg once daily with water and swallowed whole. The dose may be lowered to 2 mg daily based on individual response and tolerability; however, limited efficacy data are available for tolterodine tartrate extended-release capsules 2 mg [see **CLINICAL STUDIES** (14)].

2.2 Dosage Adjustment in Specific Populations

For patients with mild to moderate hepatic impairment (Child-Pugh Class A or B) or severe renal impairment (CrCl 10 to 30 mL/min), the recommended dose of tolterodine tartrate extended-release capsules is 2 mg once daily. Tolterodine tartrate extended-release capsules are not recommended for use in patients with severe hepatic impairment (Child-Pugh Class C). Patients with CrCl<10 mL/min have not been studied and use of tolterodine tartrate extended-release capsules in this population is not recommended [see **WARNINGS AND PRECAUTIONS** (5.6) and **USE IN SPECIFIC POPULATIONS** (8.6, 8.7)].

2.3 Dosage Adjustment in Presence of Concomitant Drugs

For patients who are taking drugs that are potent inhibitors of CYP3A4 [e.g., ketoconazole, clarithromycin, ritonavir], the recommended dose of tolterodine tartrate extended-release capsules is 2 mg once daily [see **DRUG INTERACTIONS** (7.2)].

3 DOSAGE FORMS AND STRENGTHS

The 2 mg capsules are size "4", hard gelatin capsules having a green cap and green body, imprinted "1189" on the cap with white ink, containing off-white to brownish colored pellets. The 4 mg capsules are size "3", hard gelatin capsules having a blue cap and blue body, imprinted "1190" on the cap with white ink, containing off-white to brownish colored pellets.

4 CONTRAINDICATIONS

Tolterodine tartrate extended-release capsules are contraindicated in patients with urinary retention, gastric retention, or uncontrolled narrow-angle glaucoma. Tolterodine tartrate extended-release capsules are also contraindicated in patients with known hypersensitivity to the drug or its ingredients, or to fesoterodine fumarate extended-release tablets which, like tolterodine tartrate extended-release capsules, are metabolized to 5-hydroxymethyl tolterodine [see **WARNINGS AND PRECAUTIONS** (5.2), (5.3), (5.4)].

5 WARNINGS AND PRECAUTIONS

5.1 Angioedema

Anaphylaxis and angioedema requiring hospitalization and emergency medical treatment have occurred with the first or subsequent doses of tolterodine tartrate extended-release capsules. In the event of difficulty in breathing, upper airway obstruction, or fall in blood pressure, tolterodine tartrate extended-release capsules should be discontinued and appropriate therapy promptly provided.

5.2 Urinary Retention

Administer tolterodine tartrate extended-release capsules with caution to patients with clinically significant bladder outflow obstruction because of the risk of urinary retention [see **CONTRAINDICATIONS** (4)].

5.3 Gastrointestinal Disorders

Administer tolterodine tartrate extended-release capsules with caution in patients with gastrointestinal obstructive disorders because of the risk of gastric retention. Tolterodine tartrate extended-release capsules, like other antimuscarinic drugs, may decrease gastrointestinal motility and should be used with caution in patients with conditions associated with decreased gastrointestinal motility (e.g., intestinal atony) [see **CONTRAINDICATIONS** (4)].

5.4 Controlled Narrow-Angle Glaucoma

A potent inhibitor of CYP2D6 activity, significantly inhibited the metabolism of tolterodine immediate release in CYP2D6 extensive metabolizers, resulting in a 4.8-fold increase in tolterodine AUC. There was a 52% decrease in C_{max} and a 20% decrease in AUC of 5-hydroxymethyl tolterodine (5-HMT), the pharmacologically active metabolite of tolterodine [see **CLINICAL PHARMACOLOGY** (12.1)]. The sums of unbound serum concentrations of tolterodine and 5-HMT are only 25% higher during the interaction. No dose adjustment is required when tolterodine and fluoxetine are co-administered [see **CLINICAL PHARMACOLOGY** (12.3)].

5.5 Central Nervous System Effects

Tolterodine tartrate extended-release capsules are associated with anticholinergic central nervous system (CNS) effects [see Adverse Reactions (6.2)] including dizziness and somnolence [see Adverse Reactions (6.1)]. Patients should be monitored for signs of anticholinergic CNS effects, particularly after beginning treatment or increasing the dose. Advise patients not to drive or operate heavy machinery until the drug's effects have been determined. If a patient experiences anticholinergic CNS effects, dose reduction or drug discontinuation should be considered.

5.6 Hepatic Impairment

The clearance of orally administered tolterodine immediate release was substantially lower in cirrhotic patients than in the healthy volunteers. For patients with mild to moderate hepatic impairment (Child-Pugh Class A or B), the recommended dose for tolterodine tartrate extended-release capsules is 2 mg once daily. Tolterodine tartrate extended-release capsules are not recommended for use in patients with severe hepatic impairment (Child-Pugh Class C) [see **DOSAGE AND ADMINISTRATION** (2.2) and **USE IN SPECIFIC POPULATIONS** (8.6)].

5.7 Renal Impairment

Renal impairment can significantly alter the disposition of tolterodine and its metabolites. The dose of tolterodine tartrate extended-release capsules should be reduced to 2 mg once daily in patients with severe renal impairment (CrCl: 10 to 30 mL/min). Patients with CrCl<10 mL/min have not been studied and use of tolterodine tartrate extended-release capsules in this population is not recommended [see **DOSAGE AND ADMINISTRATION** (2.2) and **USE IN SPECIFIC POPULATIONS** (8.7)].

5.8 Myasthenia Gravis

Administer tolterodine tartrate extended-release capsules with caution in patients with myasthenia gravis, a disease characterized by decreased cholinergic activity at the neuromuscular junction.

5.9 Use in Patients with Congenital or Acquired QT Prolongation

In a study of the effect of tolterodine immediate release tablets on the QT interval [see **CLINICAL PHARMACOLOGY** (12.2)], the effect on the QT interval appeared greater for 8 mg/day (two times the therapeutic dose) compared to 4 mg/day and was more pronounced in CYP2D6 poor metabolizers (PM) than extensive metabolizers (EMs). The effect of tolterodine 8 mg/day was not as large as that observed after four days of therapeutic dosing with the active control moxifloxacin. However, the confidence intervals overlapped.

These observations should be considered in clinical decisions to prescribe tolterodine tartrate extended-release capsules to patients with a known history of QT prolongation or to patients who are taking Class IA (e.g., quinidine, procainamide) or Class III (e.g., amiodarone, sotalol) antiarrhythmic medications. There has been no association of Torsade de Pointes in the international post-marketing experience with tolterodine tartrate immediate release tablets or tolterodine tartrate extended-release capsules.

6 ADVERSE REACTIONS

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

6.1 Clinical Trials Experience

The efficacy and safety of tolterodine tartrate extended-release capsules was evaluated in 1,073 patients (537 assigned to tolterodine tartrate extended-release capsules; 536 assigned to placebo) who were treated with 2, 4, 6, or 8 mg/day for up to 15 months. These included a total of 1,012 patients (505 randomized to tolterodine tartrate extended-release capsules 4 mg once daily and 507 randomized to placebo) enrolled in a randomized, placebo-controlled, double-blind, 12-week clinical efficacy and safety study.

Adverse events were reported in 52% (n=263) of patients receiving tolterodine tartrate extended-release capsules and in 49% (n=247) of patients receiving placebo. The most common adverse events reported by patients receiving tolterodine tartrate extended-release capsules were dry mouth, headache, constipation, and abdominal pain. Dry mouth was the

most frequently reported adverse event for patients treated with tolterodine tartrate extended-release capsules, occurring in 23.4% of patients treated with tolterodine tartrate extended-release capsules and 17.7% of placebo-treated patients. Dry mouth, constipation, abnormal vision (accommodation abnormalities), urinary retention, and dry eyes are expected side effects of antimuscarinic agents. A serious adverse event was reported by 1.4% (n=7) of patients receiving tolterodine tartrate extended-release capsules and by 3.6% (n=18) of patients receiving placebo.

Table 1 lists the adverse events, regardless of causality, that were reported in the randomized, double-blind, placebo-controlled 12-week study at an incidence greater than placebo and in greater than or equal to 1% of patients treated with tolterodine tartrate extended-release capsules 4 mg once daily.

Table 1. Incidence (%) of Adverse Events Exceeding Placebo Rate and Reported in ≥1% of Patients Treated with Tolterodine Tartrate Extended-Release Capsules (4 mg daily) in a 12-week, Phase 3 Clinical Trial

Body System	Adverse Event	% Tolterodine Tartrate Extended-Release Capsules n=505	% Placebo n=507
Autonomic Nervous	dry mouth	23	8
General	headache	6	5
	fatigue	2	1
Central/Peripheral Nervous	dizziness	2	1
Gastrointestinal	constipation	6	4
	abdominal pain	4	2
	dyspepsia	3	1
Vision	xerophthalmia	3	2
	vision abnormal	1	0
Psychiatric	somnolence	3	2
	anxiety	1	0
Respiratory	sinusitis	2	1
Urinary	dysuria	1	0

* in nearest integer.

The frequency of discontinuation due to adverse events was highest during the first 4 weeks of treatment. Similar percentages of patients treated with tolterodine tartrate extended-release capsules or placebo discontinued treatment due to adverse events. Dry mouth was the most common adverse event leading to treatment discontinuation among patients receiving tolterodine tartrate extended-release capsules [n=12 (2.4%) vs. placebo n=6 (1.2%)].

6.2 Post-marketing Experience

The following events have been reported in association with tolterodine use in worldwide post-marketing experience:

General: anaphylaxis and angioedema; *Cardiovascular:* tachycardia, palpitations, peripheral edema; *Gastrointestinal:* diarrhea; *Central/Peripheral Nervous:* confusion, disorientation, memory impairment, hallucinations.

Reports of aggravation of symptoms of dementia (e.g., confusion, disorientation, delusion) have been reported after tolterodine therapy was initiated in patients taking cholinesterase inhibitors for the treatment of dementia.

Because these spontaneously reported events are from the worldwide post-marketing experience, the frequency of events and the role of tolterodine in their causation cannot be reliably determined.

7 DRUG INTERACTIONS

7.1 Potent CYP2D6 Inhibitors

Fluoxetine, a potent inhibitor of CYP2D6 activity, significantly inhibited the metabolism of tolterodine immediate release in CYP2D6 extensive metabolizers, resulting in a 4.8-fold increase in tolterodine AUC. There was a 52% decrease in C_{max} and a 20% decrease in AUC of 5-hydroxymethyl tolterodine (5-HMT), the pharmacologically active metabolite of tolterodine [see **CLINICAL PHARMACOLOGY** (12.1)]. The sums of unbound serum concentrations of tolterodine and 5-HMT are only 25% higher during the interaction. No dose adjustment is required when tolterodine and fluoxetine are co-administered [see **CLINICAL PHARMACOLOGY** (12.3)].

7.2 Potent CYP3A4 Inhibitors

Ketoconazole (200 mg daily), a potent CYP3A4 inhibitor, increased the mean C_{max} and AUC of tolterodine by 2- and 2.5-fold, respectively, in CYP2D6 poor metabolizers.

For patients receiving ketoconazole or other potent CYP3A4 inhibitors such as itraconazole, clarithromycin, or ritonavir, the recommended dose of tolterodine tartrate extended-release capsules is 2 mg once daily [see **DOSAGE AND ADMINISTRATION** (2.2) and **CLINICAL PHARMACOLOGY** (12.3)].

7.3 Other Interactions

No clinically relevant interactions have been observed when tolterodine was co-administered with warfarin, with a combined oral contraceptive drug containing ethinyl estradiol and levonorgestrel, or with diuretics [see **CLINICAL PHARMACOLOGY** (12.3)].

7.4 Other Drugs Metabolized by Cytochrome P450 Isoenzymes

In vivo drug-interaction data show that tolterodine immediate release does not result in clinically relevant inhibition of CYP1A2, 2D6, 2C9, 2C19, or 3A4 as evidenced by lack of influence on the marker drugs caffeine, debrisoquine, 5-warfarin, and omeprazole [see **CLINICAL PHARMACOLOGY** (12.3)].

7.5 Drug-Laboratory-Test Interactions

Interactions between tolterodine and laboratory tests have not been studied.

7.6 Other Anticholinergics

The concomitant use of tolterodine tartrate extended-release capsules with other anticholinergic (antimuscarinic) agents may increase the frequency and/or severity of dry mouth, constipation, blurred vision, somnolence, and other anticholinergic pharmacological effects.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary
There are no available data with tolterodine tartrate extended-release capsules use in pregnant women to inform drug-associated risks. In animal reproduction studies, oral administration of tolterodine and its 5-HMT metabolite to pregnant mice during organogenesis did not produce adverse developmental outcomes at doses approximately 9 to 12 times the clinical exposure at a dose of 20 mg/kg/day; however, higher doses produced adverse developmental outcomes [see **Data**].

In the U.S. general population, the estimated background rate of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Animal Data

No anomalies or malformations were observed after oral administration of tolterodine to pregnant mice during organogenesis at approximately 9 to 12 times the clinical exposure to the pharmacologically active components of tolterodine tartrate extended-release capsules (based on the AUC of tolterodine and its 5-HMT metabolite at a dose of 20 mg/kg/day). At 14 to 18 times the clinical exposure (doses of 30 to 40 mg/kg/day) in mice, tolterodine was embryo-lethal, caused reduced fetal weight, and increased the incidence of fetal abnormalities (cleft palate, digital abnormalities, intra-abdominal hemorrhage, and various skeletal abnormalities, primarily reduced ossification). Pregnant rabbits administered tolterodine subcutaneously at about 0.3 to 2.5 times the clinical exposure (dose of 0.8 mg/kg/day) did not show any embryotoxicity or teratogenicity.

8.2 Lactation

Risk Summary
There is no information on the presence of tolterodine or its 5-HMT metabolite in human milk, the effects on the breastfed infant, or the effects on milk production. Based on limited data, tolterodine is excreted into the milk in mice in low amounts [see **Data**]. The development and health benefits of breastfeeding should be considered along with the mother's clinical need for tolterodine tartrate extended-release capsules and any potential adverse effects on the breastfed infant from tolterodine tartrate extended-release capsules or from the underlying maternal condition.

There is no information on the presence of tolterodine or its 5-HMT metabolite in human milk, the effects on the breastfed infant, or the effects on milk production. Based on limited data, tolterodine is excreted into the milk in mice in low amounts [see **Data**]. The development and health benefits of breastfeeding should be considered along with the mother's clinical need for tolterodine tartrate extended-release capsules and any potential adverse effects on the breastfed infant from tolterodine tartrate extended-release capsules or from the underlying maternal condition.

Animal Data
The use of radiolabeled tolterodine in pregnant mice produced milk: plasma ratios that ranged between 0.0 and 0.7.

8.4 Pediatric Use

The effectiveness of tolterodine tartrate extended-release capsules has not been established in pediatric patients.

Efficacy was not established in two randomized, placebo-controlled, double-blind, 12-week studies that enrolled 710 pediatric patients (486 on tolterodine tartrate extended-release capsules, 224 on placebo) aged 5 to 10 years with urinary frequency and urge incontinence. The percentage of patients with urinary tract infections was higher in patients treated with tolterodine tartrate extended-release capsules (6.6%) compared to patients who received placebo (4.5%). Aggressive, abnormal, and hyperactive behavior and attention disorders occurred in 2.9% of children treated with tolterodine tartrate extended-release capsules compared to 0.9% of children treated with placebo.

8.5 Geriatric Use

No overall differences in safety were observed between the older and younger patients treated with tolterodine.

In multiple-dose studies in which tolterodine immediate release 4 mg (2 mg bid) was administered

PRODUCT NAME :	Tolterodine Tartrate Extended-Release Capsules	COUNTRY : US	LOCATION : Indrad	Supersedes A/W No.:	8102981	V. No.: 01	
ITEM / PACK :	Outsert	NO. OF COLORS: 1	REMARK :				
DESIGN STYLE :	Back	PANTONE SHADE NOS.:	SUBSTRATE : 40 g/m ² Bible Paper				
CODE :	8106403	Black	Activities	Department	Name	Signature	Date
DIMENSIONS (MM) :	490 x 340		Prepared By	Pkg.Dev			
ART WORK SIZE :	S/S		Reviewed By	Pkg.Dev			
DATE :	03-01-2026		Reviewed By	Quality			
		Font Size 6.5 pt_Medi Font Size 10 pt	Approved By	Quality			

**Note: Pharma code/ Bar code and adjacent text must be visible on folded leaflet.
These details can be moved by printed to arrange pharma code/ Bar code and adjacent text visible on folded leaflet.**

Drug Interactions:

Potent CYP2D6 inhibitors: Fluoxetine is a selective serotonin reuptake inhibitor and a potent inhibitor of CYP2D6 activity. In a study to assess the effect of fluoxetine on the pharmacokinetics of tolterodine immediate release and its metabolites, it was observed that fluoxetine significantly inhibited the metabolism of tolterodine immediate release in extensive metabolizers, resulting in a 4.8-fold increase in tolterodine AUC. There was a 52% decrease in C_{max} and a 20% decrease in AUC of 5-hydroxymethyl tolterodine (5-HMT, the pharmacologically active metabolite of tolterodine). Fluoxetine thus alters the pharmacokinetics in patients who would otherwise be CYP2D6 extensive metabolizers of tolterodine immediate release to resemble the pharmacokinetic profile in poor metabolizers. The sums of unbound serum concentrations of tolterodine immediate release and 5-HMT are only 25% higher during the interaction. No dose adjustment is required when tolterodine and fluoxetine are co-administered.

Potent CYP3A4 inhibitors: The effect of a 200 mg daily dose of ketoconazole on the pharmacokinetics of tolterodine immediate release was studied in 8 healthy volunteers, all of whom were CYP2D6 poor metabolizers. In the presence of ketoconazole, the mean C_{max} and AUC of tolterodine increased by 2- and 2.5-fold, respectively. Based on these findings, other potent CYP3A4 inhibitors may also lead to increases of tolterodine plasma concentrations.

For patients receiving ketoconazole or other potent CYP3A4 inhibitors such as itraconazole, miconazole, clarithromycin, ritonavir, the recommended dose of tolterodine tartrate extended-release capsules is 2 mg daily [see **DOSE AND ADMINISTRATION (2.3)**].

Warfarin: In healthy volunteers, coadministration of tolterodine immediate release 4 mg (2 mg bid) for 7 days and a single dose of warfarin 25 mg on day 4 had no effect on prothrombin time, Factor VII suppression, or on the pharmacokinetics of warfarin.

Oral Contraceptives: Tolterodine immediate release 4 mg (2 mg bid) had no effect on the pharmacokinetics of an oral contraceptive (ethinyl estradiol 30 mcg/levonorgestrel 150 mcg) as evidenced by the monitoring of ethinyl estradiol and levonorgestrel over a 2-month period in healthy female volunteers.

Diuretics: Coadministration of tolterodine immediate release up to 8 mg (4 mg bid) for up to 12 weeks with diuretic agents, such as indapamide, hydrochlorothiazide, triamterene, bendroflumethiazide, chlorothiazide, methylchlorothiazide, or furosemide, did not cause any adverse electrocardiographic (ECG) effects.

Effect of tolterodine on other drugs metabolized by Cytochrome P450 enzymes: Tolterodine immediate release does not cause clinically significant interactions with other drugs metabolized by the major drug-metabolizing CYP enzymes. *In vivo* drug-interaction data show that tolterodine immediate release does not result in clinically relevant inhibition of CYP1A2, 2D6, 2C9, 2C19, or 3A4 as evidenced by lack of influence on the marker drugs caffeine, debrisoquine, S-warfarin, and omeprazole. *In vitro* data show that tolterodine immediate release is a competitive inhibitor of CYP2D6 at high concentrations (K_i 1.05 μM), while tolterodine immediate release as well as the 5-HMT are devoid of any significant inhibitory potential regarding the other isoenzymes.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity studies with tolterodine were conducted in mice and rats. At the maximum tolerated dose in mice (30 mg/kg/day), female rats (20 mg/kg/day), and male rats (30 mg/kg/day), exposure margins were approximately 6 to 9 times, 7 times, and 11 times the clinical exposure to the pharmacologically active components of tolterodine tartrate extended-release capsules (based on AUC of tolterodine and its 5-HMT metabolite). At these exposure margins, no increase in tumors was found in either mice or rats.

No mutagenic or genotoxic effects of tolterodine were detected in a battery of *in vitro* tests, including bacterial mutation assays (Ames test) in 4 strains of *Salmonella typhimurium* and in 2 strains of *Escherichia coli*; a gene mutation assay in L5178Y mouse lymphoma cells; and chromosomal aberration tests in human lymphocytes. Tolterodine was also negative *in vivo* in the bone marrow micronucleus test in the mouse.

In female mice treated for 2 weeks before mating and during gestation with 20 mg/kg/day (about 9 to 12 times the clinical exposure via AUC), neither effects on reproductive performance or fertility were seen. In male mice, a dose of 30 mg/kg/day did not induce any adverse effects on fertility.

14 CLINICAL STUDIES

Tolterodine tartrate extended-release capsules 2 mg were evaluated in 29 patients in a Phase 2 dose-effect study. Tolterodine tartrate extended-release capsules 4 mg were evaluated for the treatment of overactive bladder with symptoms of urge urinary incontinence and frequency in a randomized, placebo-controlled, multicenter, double-blind, Phase 3, 12-week study. A total of 507 patients received tolterodine tartrate extended-release capsules 4 mg once daily in the morning and 508 received placebo. The majority of patients were Caucasian (95% and female (81%), with a mean age of 61 years (range, 20 to 93 years). In the study, 642 patients (42%) were 65 to 93 years of age. The study included patients known to be responsive to tolterodine immediate release and other anticholinergic medications, however, 47% of patients never received prior pharmacotherapy for overactive bladder. At study entry, 97% of patients had at least 5 urge incontinence episodes per week and 91% of patients had 8 or more micturitions per day.

The primary efficacy assessment was change in mean number of incontinence episodes per week at week 12 from baseline. Secondary efficacy measures included change in mean number of micturitions per day and mean volume voided per micturition at week 12 from baseline.

Patients treated with tolterodine tartrate extended-release capsules experienced a statistically significant decrease in number of urinary incontinence per week from baseline to last assessment (week 12) compared with placebo as well as a decrease in the average daily urinary frequency and an increase in the average urine volume per void.

Mean change from baseline in weekly incontinence episodes, urinary frequency, and volume voided between placebo and tolterodine tartrate extended-release capsules are summarized in Table 4.

Table 4. 95% Confidence Intervals (CI) for the Difference between Tolterodine Tartrate Extended-Release Capsules (4 mg daily) and Placebo for Mean Change at Week 12 from Baseline*

	Tolterodine Tartrate Extended-Release Capsules (n=507)	Placebo (n=508) [†]	Treatment Difference, vs. Placebo (95% CI)
Number of incontinence episodes/week			
Mean Baseline	22.1	23.3	-4.8 [‡]
Mean Change from Baseline	-11.8 (SD 17.8)	-6.9 (SD 15.4)	(-6.9, -2.8)
Number of micturitions/day			
Mean Baseline	10.9	11.3	-0.6 [‡]
Mean Change from Baseline	-1.8 (SD 3.4)	-1.2 (SD 2.9)	(-1.0, -0.2)
Volume voided per micturition (mL)			
Mean Baseline	141	136	20 [‡]
Mean Change from Baseline	34 (SD 51)	14 (SD 41)	(14, 26)

SD = Standard Deviation.

* Intent-to-treat analysis.

[†] 1 to 2 patients missing in placebo group for each efficacy parameter.

[‡] The difference between tolterodine tartrate extended-release capsules and placebo was statistically significant.

16 HOW SUPPLIED/STORAGE AND HANDLING

Tolterodine tartrate extended-release capsules, USP are supplied as follows:

2 mg capsules are size "4", hard gelatin capsules having a green cap and green body, imprinted "1189" on the cap with white ink, containing off-white to brownish colored pellets.

Bottles of 30	NDC 13668-189-30
Bottles of 90	NDC 13668-189-90
Bottles of 500	NDC 13668-189-05
100 Unit dose capsules	NDC 13668-189-74

4 mg capsules are size "3", hard gelatin capsules having a blue cap and blue body, imprinted "1190" on the cap with white ink, containing off-white to brownish colored pellets.

Bottles of 30	NDC 13668-190-30
Bottles of 90	NDC 13668-190-90
Bottles of 500	NDC 13668-190-05
100 Unit dose capsules	NDC 13668-190-74

Store tolterodine tartrate extended-release capsules, USP at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Protect from light.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Antimuscarinic Effects

Inform patients that antimuscarinic agents such as tolterodine tartrate extended-release capsules may have side effects including blurred vision, dizziness, or drowsiness. Advise patients not to drive, operate machinery, or do other potentially dangerous activities until they know how tolterodine tartrate extended-release capsules affects them.

PATIENT INFORMATION

Tolterodine Tartrate (tol-TER-oh-deen TAR-trate) Extended-Release Capsules, USP

Read the Patient Information that comes with tolterodine tartrate extended-release capsules before you start using them and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your doctor about your condition or your treatment. Only your doctor can determine if treatment with tolterodine tartrate extended-release capsules is right for you.

What are tolterodine tartrate extended-release capsules?

Tolterodine tartrate extended-release capsules are a prescription medicine for **adults** used to treat the following symptoms due to a condition called **overactive bladder**:

- Having a strong need to urinate with leaking or wetting accidents (urge urinary incontinence).
- Having a strong need to urinate right away (urgency).
- Having to urinate often (frequency).

Tolterodine tartrate extended-release capsules did not help the symptoms of overactive bladder when studied in children.

What is overactive bladder?

Overactive bladder happens when you cannot control your bladder muscle. When the muscle contracts too often or cannot be controlled, you get symptoms of overactive bladder, which are leakage of urine (urge urinary incontinence), needing to urinate right away (urgency), and needing to urinate often (frequency).

Who should not take tolterodine tartrate extended-release capsules?

Do not take tolterodine tartrate extended-release capsules if:

- You have trouble emptying your bladder (also called “urinary retention”).
- Your stomach empties slowly (also called “gastric retention”).
- You have an eye problem called “uncontrolled narrow-angle glaucoma”.
- You are allergic to tolterodine tartrate extended-release capsules or to any of its ingredients. See the end of this leaflet for a complete list of ingredients.
- You are allergic to Toviaz[®], which contains fesoterodine.

What should I tell my doctor before starting tolterodine tartrate extended-release capsules?

Before starting tolterodine tartrate extended-release capsules, tell your doctor about all of your medical conditions, including if you:

- Have any stomach or intestinal problems.
- Have trouble emptying your bladder or you have a weak urine stream.
- Have an eye problem called narrow-angle glaucoma.
- Have liver problems.
- Have kidney problems.
- Have a condition called myasthenia gravis.
- Or any family members have a rare heart condition called QT prolongation (long QT syndrome).
- Are pregnant or trying to become pregnant. It is not known if tolterodine tartrate extended-release capsules could harm your unborn baby.
- Are breastfeeding. It is not known if tolterodine tartrate passes into your milk and if it can harm your child.

Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. Other drugs can affect how your body handles tolterodine tartrate extended-release capsules. Your

doctor may use a lower dose of tolterodine tartrate extended-release capsules if you are taking:

- Certain medicines for fungus or yeast infections such as Nizoral[®] (ketoconazole), Sporanox[®] (itraconazole), or Monistat[®] (miconazole).
- Certain medicines for bacteria infections such as Biaxin[®] (clarithromycin).
- Certain medicines for treatment of HIV infection such as Norvir[®] (ritonavir), Invirase[®] (saquinavir), Reyataz[®] (atazanavir).
- Sandimmune[®] (cyclosporine) or Velban[®] (vinblastine).

Know the medicines you take. Keep a list of them with you to show your doctor or pharmacist each time you get a new medicine.

How should I take tolterodine tartrate extended-release capsules?

- Take tolterodine tartrate extended-release capsules exactly as prescribed. Your doctor will prescribe the dose that is right for you. Do not change your dose unless told to do so by your doctor.
- Take tolterodine tartrate extended-release capsules once a day with liquid. Swallow the whole capsule. Tell your doctor if you cannot swallow a capsule.
- Tolterodine tartrate extended-release capsules can be taken with or without food.
- Take tolterodine tartrate extended-release capsules the same time each day.
- If you miss a dose of tolterodine tartrate extended-release capsules, begin taking tolterodine tartrate extended-release capsules again the next day. Do not take 2 doses of tolterodine tartrate extended-release capsules in the same day.
- If you took more than your prescribed dose of tolterodine tartrate extended-release capsules, call your doctor or poison control center, or go to the hospital emergency room.

What are possible side effects of tolterodine tartrate extended-release capsules?

Tolterodine tartrate extended-release capsules may cause allergic reactions that may be serious. Symptoms of a serious allergic reaction may include swelling of the face, lips, throat, or tongue. If you experience these symptoms, you should stop taking tolterodine tartrate extended-release capsules and get emergency medical help right away.

The most common side effects with tolterodine tartrate extended-release capsules are:

- Dry mouth
- Headache
- Constipation
- Stomach pain

Medicines like tolterodine tartrate extended-release capsules can cause blurred vision, dizziness, and drowsiness.

Do not drive, operate machinery, or do other dangerous activities until you know how tolterodine tartrate extended-release capsules affect you.

Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088.

These are not all the side effects with tolterodine tartrate extended-release capsules. For a complete list, ask your doctor or pharmacist.

How do I store tolterodine tartrate extended-release capsules?

- Store tolterodine tartrate extended-release capsules at room temperature, 68° to 77°F (20° to 25°C); brief periods permitted between 59° to 86°F (15° to 30°C). Protect from light. Keep in a dry place.
- **Keep tolterodine tartrate extended-release capsules and all medicines out of the reach of children.**

General Information about tolterodine tartrate extended-release capsules

Medicines are sometimes prescribed for conditions that are not in the patient information leaflet. Only use tolterodine tartrate extended-release capsules the way your doctor tells you. Do not share it with other people even if they have the same symptoms you have. It may harm them.

This leaflet summarizes the most important information about tolterodine tartrate extended-release capsules. If you

would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about tolterodine tartrate extended-release capsules that is written for health professionals. You can also call 1-800-912-9561.

What are the ingredients in tolterodine tartrate extended-release capsules?

Active ingredients: tolterodine tartrate, USP

Inactive ingredients: cellulose microcrystalline, ethyl cellulose, FD & C blue 2, gelatin, hypromellose, medium chain triglycerides, oleic acid, sucrose, talc and titanium dioxide. 2 mg capsule also contains iron oxide yellow. Capsules have pharmaceutical grade printing ink that contains potassium hydroxide, propylene glycol, shellac, strong ammonia solution and titanium dioxide.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Trademarks are the property of their respective owners.

TORRENT[®] PHARMA

Manufactured by:
Torrent Pharmaceuticals LTD., India.

Manufactured for:
Torrent Pharma INC., Basking Ridge, NJ 07920.

8106403

Revised: January 2026