

VIZYLAC DT

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory Only

Abbreviated Prescribing information for VIZYLAC DT [Lactic Acid Bacillus and Zinc Sulphate Monohydrate Tablets]

[Please refer the complete prescribing information available at www.torrentpharma.com]

PHARMACOLOGICAL PROPERTIES:

MECHANISM OF ACTION: Zinc: The physiological effect of zinc on intestinal ion transport has not yet been established thoroughly. Lactic Acid Bacillus: The mechanism of action is presumed to be a result of improving gastrointestinal ecology made possible by replenishing the quantity of desirable obligate microorganisms and antagonizing pathogenic microbes.

INDICATIONS: It is indicated for the treatment of Diarrhea.

DOSAGE AND ADMINISTRATION: As directed by the physician. Disperse the tablet in a teaspoonful of previously boiled and cooled water before administration.

CONTRAINDICATION: Hypersensitivity to Lactic Acid Bacillus and Zinc sulphate and other ingredients listed in prescribing information. Copper deficiency.

WARNINGS & PRECAUTIONS: Accumulation of zinc may occur in cases of renal failure. This product contains sodium. This should be taken into consideration by patients on a controlled sodium diet. Probiotics do not appear to pose any safety concerns for pregnant and lactating women. Systemic absorption is rare when probiotics are used by healthy individuals, and the current literature does not indicate an increase in adverse pregnancy outcomes.

DRUG INTERACTIONS: Zinc may inhibit the absorption of copper. Zinc may reduce the absorption of concurrently administered tetracycline, also the absorption of zinc may be reduced by tetracycline; when both are being given an interval of at least three hours should be allowed. Zinc may reduce the absorption of quinolones; ciprofloxacin, levofloxacin, moxifloxacin, norfloxacin and ofloxacin. The absorption of zinc may be reduced by calcium salts. The absorption of zinc may be reduced by oral iron, also the absorption of oral iron may be reduced by zinc. The absorption of zinc may be reduced by Penicillamine, also the absorption of penicillamine may be reduced by zinc. The absorption of zinc may be reduced by trientine, also the absorption of trientine may be reduced by zinc.

ADVERSE REACTIONS: Abdominal detention, Bloating, Skin rashes, Loss of appetite, Altered sleep patterns, dyspepsia, nausea, vomiting, diarrhoea, gastric irritation, gastritis, irritability, headache, lethargy.

MARKETED BY:

TORRENT
PHARMA

Torrent Pharmaceuticals Limited.

IN/VIZYLAC DT 120x10⁶, 10 mg/APR-2026/03/ABPI

(Additional information is available on request)