

## TRUECEPROL

**For the use of a Registered Medical Practitioner or a Hospital or a Laboratory Only**

Abbreviated Prescribing information for Trueceprol [Oxaceprol Capsules 200mg] [Please refer the complete prescribing information available at [www.torrentpharma.com](http://www.torrentpharma.com)]

### PHARMACOLOGICAL PROPERTIES:

**MECHANISM OF ACTION:** The exact mechanism of action of oxaceprol is not known. It affects connective tissue metabolism and has been used in dermatology, to promote wound healing, and in rheumatic disorders. Experimental models of arthritis showed that oxaceprol reduced leukocyte extravasation, as well as the adhesion of leukocytes to capillaries. In vitro studies have shown that oxaceprol stimulates the uptake of <sup>3</sup>H-glucosamine and <sup>3</sup>H-proline in chondrocytes and the incorporation of <sup>3</sup>H-proline in the macromolecular structure of the matrix of cartilage; which would be responsible for the activity on regenerating tissue.

**INDICATION:** For the relief of sign and symptoms of osteoarthritis in adult patients.

**DOSAGE AND ADMINISTRATION:** Oxaceprol hard gelatin capsule are preferably taken before a meal, unchewed, with sufficient liquid. The period of time for which the drug is to be administered depends on the character and intensity of the affection and is to be fixed individually. **Adult:** 200 mg tid. In severe cases, initial dose can be increased to 400 mg tid. Taken before a meal, unchewed.

**Elderly:** Dose reduction is recommended.

**CONTRAINDICATION:** A known hypersensitivity to Oxaceprol or one of the other components.

**WARNINGS & PRECAUTIONS:** Special precautions should be taken while use of oxaceprol in children, elderly, pregnancy and lactation.

**DRUG INTERACTIONS:** Affects coagulation in patients under anticoagulative therapy with Vitamin K-antagonists. A control of the prothrombin time under simultaneous therapy with oxaceprol is, therefore, recommended. It may decrease effects of anti-hypertensives and also affects toxicity of methotrexate and lithium.

**ADVERSE REACTIONS:** Under the treatment with Oxaceprol occasionally there have been observed gastrointestinal complaints, such as nausea, impaired appetite, pain in the stomach or diarrhea, which are often of a passing nature. Seldom there may occur allergic reactions (a reddening of the skin, an itching of the skin, exanthema). In exceptional cases the following reactions of allergic genesis were described: loss of hair, arthralgia, vasculitis, urticaria, giant urticaria and allergic eosinophilia.

**MARKETED BY:**

**TORRENT**  
PHARMA

TORRENT PHARMACEUTICALS LTD.

**IN/TRUECEPROL 200mg/MAR 2026/03/ABPI**

(Additional information is available on request)